

Clauna-Lumanta, M. M. T., et al. (2016). "The effects of GCSF on the recovery time and duration of hospitalization in patients with anti-thyroid drug-induced agranulocytosis in a tertiary hospital." *Journal of the ASEAN Federation of Endocrine Societies* 31(2): 131-136.

Concepcion, J., et al. (2015). "Rapid recovery of ANC after using GCSF to treat dose-related methimazole-induced agranulocytosis in a pediatric graves' disease patient." *Endocrine Reviews* 36.

Garla, V. V. and A. Subauste (2014). "Thyroid storm and methimazole induced agranulocytosis in a patient: A treatment conundrum." *Endocrine Reviews* 35.

IonicĂ, M., et al. (2014). "Severe and prolonged febrile agranulocytosis under thyrosol." *BMC Infectious Diseases* 14(7).

Jain, M., et al. (2013). "Methimazole-induced concomitant agranulocytosis and hepatotoxicity." *Endocrine Reviews* 34(3).

Kim, H. K., et al. (2015). "Methimazole is associated with more severe agranulocytosis: Multicenter study in Korea." *Endocrine Reviews* 36.

Majors, C., et al. (2016). "Agranulocytosis after treatment with methimazole." *Endocrine Reviews* 37(2).

Mendes, P. and R. Carvalho (2001). "Diabetes mellitus and hiperthyroidism." *Arquivos de Medicina* 15(4-6): 81-85.

Minamitani, K., et al. (2011). "A report of three girls with antithyroid drug-induced agranulocytosis; retrospective analysis of 18 cases aged 15 years or younger reported between 1995 and 2009." *Clinical Pediatric Endocrinology* 20(2): 39-46.

Yang, J., et al. (2016). "Characteristics of Antithyroid Drug-Induced Agranulocytosis in Patients with Hyperthyroidism: A Retrospective Analysis of 114 Cases in a Single Institution in China Involving 9690 Patients Referred for Radioiodine Treatment over 15 Years." *Thyroid* 26(5): 627-633.

Zelada, H., et al. (2016). "In-hospital mortality secondary to methimazole-induced agranulocytosis." *Thyroid* 26: A158-A159.